

MAR09-2008-004199

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Sorting Category: 23.12.3 (E)

Measurement of Autler-Townes and Mollow transitions in a strongly driven superconducting qubit M. BAUR, S. FILIPP, ETH QUANTUM DEVICE TEAM, ETH Zurich, A. BLAIS, Université de Sherbrooke, A. WALLRAFF, ETH Zurich — The spectrum of a multilevel atom can be significantly modified when interacting with electromagnetic fields. In the simplest case where a two-level atom is driven on resonance, two sidebands offset from the main atomic line by the Rabi frequency Ω appear in the fluorescence spectrum, referred to as the Mollow triplet. Similarly, when probing transitions into a third atomic level, the absorption spectrum shows two spectral lines separated by Ω , called the Autler-Townes doublet. Here we present a measurement of the Autler-Townes doublet and the sidebands of the Mollow triplet in a strongly driven superconducting qubit. The corresponding transitions are detected using dispersive read-out of the qubit coupled off-resonantly to a microwave transmission line resonator. The observed frequencies of the Autler-Townes and Mollow spectral lines are in excellent agreement with a generalized dispersive Jaynes-Cummings model.

- Prefer Oral Session
 Prefer Poster Session

Matthias Baur
mabaur@phys.ethz.ch
ETH Zurich

Special instructions: membership pending

Date submitted: 21 Nov 2008

Electronic form version 1.4